



Correction: Effects of isotretinoin on new bone formation after maxillary sutural expansion

Musa Bulut¹ · Yasemin Nur Korkmaz¹ · Sevilay Erimsah²

Published online: 9 December 2020
© Springer Medizin Verlag GmbH, ein Teil von Springer Nature 2020

Correction: Auswirkungen von Isotretinoin auf die Knochenneubildung nach maxillärer suturaler Expansion

Correction to:

J Orofac Orthop (2020) 81:440–446

<https://doi.org/10.1007/s00056-019-00209-2>

The original version of this article unfortunately contained a mistake. The third bullet under the section Materials and Methods should read “Soybean oil (SOY) group: Soybean oil was administered via oral gavage (**2 ml/kg**) during the nursery period before expansion (an interval of 40 days), and during the expansion (period of 5 days) + retention period (period of 12 days—a total of 57 days)” instead of “Soybean oil (SOY) group: Soybean oil was administered via oral gavage (0.2 ml/kg) during the nursery period before expansion (an interval of 40 days), and during the expansion (period of 5 days) + retention period (period of 12 days—a total of 57 days)”.

The authors apologize for the mistake.

Published online: 21 January 2020
© Springer Medizin Verlag GmbH, a part of Springer Nature 2019

The online version of the original article can be found under
<https://doi.org/10.1007/s00056-019-00209-2>.

✉ Musa Bulut, DDS, PhD
musabulut@gmail.com

¹ Faculty of Dentistry, Department of Orthodontics, Bolu Abant İzzet Baysal University, Gölköy Yerleşkesi, 14030 Bolu, Turkey

² Faculty of Medicine, Department of Histology and Embryology, Bolu Abant İzzet Baysal University, Bolu, Turkey