



## Evaluation of hemogram parameters in diabetic patients with coronary artery ectasia

Mehmet Inanir 

Department of Cardiology, Bolu Abant Izzet Baysal University, School of Medicine, Bolu, Turkey

### ABSTRACT

**Aim:** To compare the importance of hemogram parameters in predicting the disease in diabetic patients with coronary artery ectasia (CAE) and normal coronary artery.

**Methods:** The records of 7287 patients who underwent coronary angiography between January 2017 and October 2019 were reviewed. After appropriate exclusions, diabetic patients were divided into coronary artery ectasia and normal coronary artery groups. A total of 248 patients were included in the study and hemogram parameters of these two groups were compared.

**Results:** Compared to control group white blood count (WBC) [8 (4-13) vs. 7 (5-12)  $u/mm^3$ ,  $p=0.023$ ], hemoglobin [13 (10-16) vs. 14 (10-20)  $gr/dL$ ,  $p=0.015$ ], red cell distribution width (RDW) [16 (14-20) vs. 15 (12-19) %,  $p=0.026$ ], neutrophil [4.5 (2.1-11.4) vs. 4.0 (0.2-7.5)  $u/mm^3$ ,  $p=0.003$ ], platelet counts (Plt) [266 (196-450) vs. 236 (163-362)  $k/mm^3$   $p<0.001$ ], platelet distribution width (PDW) (17.9 (16.2-20.4) vs. 17.7 (15.9-19.7) %  $p=0.011$ ), mean platelet volume (MPV) [8.4 (6.4-11.2) vs. 7.9 (6.6-10.1)  $Fl$ ,  $p=0.015$ ], plateletcrit (PCT) [0.20 (0.14-0.32) vs. 0.19 (0.13-0.26),  $p<0.001$ ], and neutrophil lymphocyte ratio (NLR) [2.1 (1.0-9.7) vs. 1.6 (0.2-5.7),  $p=0.002$ ] were significantly higher in CAE patients.

**Conclusion:** The results of this study suggest that the increased some hemogram parameters may be useful in predicting disease in diabetic patients with CAE.

**Keywords:** Coronary artery ectasia, diabetes mellitus, hemogram parameters.

© 2020 [experimentalbiomedicalresearch.com](http://www.experimentalbiomedicalresearch.com)

✉ Dr. Mehmet Inanir,

Department of Cardiology, Bolu Abant Izzet Baysal University, School of Medicine, Bolu, Turkey

E-mail: [mdmehmetinanir@yahoo.com](mailto:mdmehmetinanir@yahoo.com)

Received: 2019-11-23 / Revisions: 2019-12-29

Accepted: 2020-01-01 / Publication Date: 2020-03-06

### Introduction

Coronary artery ectasia (CAE) is defined as the dilation of the coronary artery lumen. The term "ectasia" is defined as the widespread expansion of a coronary artery, whereas focal coronary expansion is called "coronary aneurysm" [1]. CAE is a disease of the coronary

arteries with abnormal dilatation of the coronary arteries [2]. The diameter of the dilated segment is 1.5 times the diameter of the normal adjacent segment [3]. CAE is a form of coronary atherosclerosis characterized by internal and external elastic lamina disorder [4, 5]. The incidence of CAE ranges from 0.3 to 4.990 [6, 7]. CAE can be seen due to genetics, Kawasaki disease, mycotic or septic emboli, Marfan syndrome, polyarthritis nodosa arthritis, Takayasu disease, systemic lupus erythematosus, hypertension, smoking, cocaine use, percutaneous transluminal coronary

angioplasty, stent, and directional coronary atherectomy. CAE is also related to apical hypertrophic cardiomyopathy [8].

The etiopathogenesis of CAE is not fully understood [9, 10]. Although the main cause of coronary artery ectasia is unknown, atherosclerosis has been the most accused pathogenesis [11]. One of the most important indicators of atherosclerotic processes is endothelial dysfunction [12, 13]. CAE has commonly been evaluated as a variant of atherosclerotic heart disease. However; more intense inflammation has been detected in CAE than obstructive coronary artery disease (CAD) [14]

Histology is usually due to chronic vascular inflammation that shows thickened fibrotic intima with lipid accumulation [15]. The thinning of the tunica media environment associated with chronic inflammation is considered to be the main pathogenesis of extensive remodeling [10]. Ectasia can lead to slow flow in the coronary arteries, dissection, thrombus formation, and vasospasm [16, 17]. The primary symptom of CAE is chest pain. Ectasia can cause the acute coronary syndrome, ventricular arrhythmias and sudden cardiac death without severe coronary artery stenosis [7]. Coronary angiography is the gold standard test for the diagnosis of coronary artery ectasia [8].

Diabetes mellitus (DM) is an important public health problem due to high morbidity and mortality from microvascular and macrovascular complications [18]. Endothelial dysfunction is the basis for the development of long-term complications of diabetes [19]. The activity and aggregation of platelets are important in terms of thrombus during the atherogenesis process [20, 21]. Therefore, in this study, we aimed to evaluate the hemogram parameters of diabetic CAE patients.

## Materials and Methods

We reviewed 7287 angiograms performed between January 2017 and October 2019, from Bolu Abant Izzet Baysal University Medical Faculty Hospital. "The Siemens Axiom Artis diagnostic device (Siemens Healthcare GmbH, Forchheim, Germany)" was used to perform coronary angiography. Coronary angiography (CAG) was performed to investigate ischemic heart diseases based on clinical indications. The study was conducted in accordance with the ethical approval of the University Ethics Committee. (Date: 24/10/2019; Decision number: 2019/217) Data about patients were obtained from the institution's database and patient files. CAG images recorded in digital format were evaluated visually by two blind cardiologists and patients diagnosed as CAE were included in the study. Patients included in the study were selected from patients with chronic coronary syndrome (CCS). Patients with clear CAE evidence were selected. The baseline demographic data and clinical cardiovascular risk factors; hypertension, diabetes mellitus, smoking or ex-smoking, family history of CAD, dyslipidemia weight and height were determined from hospital records. There was no significant difference in demographic parameters between CAE patients and the control group (normal coronary angiography). Subjects with a history of chronic diseases such as heart failure (ejection fraction <50%), acute coronary syndrome (ACS), previous coronary artery bypass grafting, percutaneous coronary intervention, significant valve disease, patients under 18 years of age, atrial fibrillation, hypertension, smoking, autoimmune diseases, pregnancy, iatrogenic ectasia, myocarditis, pericarditis, acute and chronic lung disease, obstructive sleep apnea, chronic inflammation, active infection, cancer, immunosuppressive therapy,

hypo/hyperthyroidism, stroke, mental retardation, delirium, dementia, any hematological abnormality (sickle cell anemia, thrombocytopenia etc.), and antiplatelet / anticoagulant agents and steroid users, liver or kidney failure and electrolyte imbalance were excluded from the study.

### Statistical analysis

Statistical analysis was conducted with SPSS software (SPSS 22.0 for Windows, IBM Co, Chicago, IL, USA). Kolmogorov Smirnov test was used to determine distribution normality. Normal variables were compared with the T-test and expressed as mean  $\pm$  standard deviation. Mann Whitney U test was used for variables showing the abnormal distribution and expressed as median (IQR: interquartile interval). A chi-square test was used for comparison of nonparametric variables. A p-value lower than 0.05 was considered statistically significant.

### Results

We enrolled 248 individuals including 124 CAE patients (mean age: 61.4 $\pm$ 10.9 years) and

124 control persons (mean age: 59.3 $\pm$ 10.3 years). The mean age was 61.4 $\pm$ 10.9 and 59.3 $\pm$ 10.3 in the patient and the control groups, respectively. Frequencies of sex, and body mass index (BMI) were not significantly different between the patient and the control groups (Table 1). Compared to control group white blood count (WBC) [8 (4-13) vs. 7 (5-12) u/mm<sup>3</sup>,  $p=0.023$ ], hemoglobin [13 (10-16) vs. 14 (10-20) gr/dL,  $p=0.015$ ], red cell distribution width (RDW) [16 (14-20) vs. 15 (12-19) %,  $p=0.026$ ], neutrophil [4.5 (2.1-11.4) vs. 4.0 (0.2-7.5) u/mm<sup>3</sup>,  $p=0.003$ ], platelet counts (Plt) [266 (196-450) vs. 236 (163-362) k/mm<sup>3</sup>  $p<0.001$ ], platelet distribution width (PDW) (17.9 (16.2-20.4) vs. 17.7 (15.9-19.7) %  $p=0.011$ ), mean platelet volume (MPV) [8.4 (6.4-11.2) vs. 7.9 (6.6-10.1) fL,  $p=0.015$ ], plateletcrit (PCT) [0.20 (0.14-0.32) vs. 0.19 (0.13-0.26),  $p=0.001$ ], and neutrophil lymphocyte ratio (NLR) [2.1 (1.0-9.7) vs. 1.6 (0.2-5.7),  $p=0.002$ ] were significantly higher in CAE patients. There was no significant difference between the two groups in terms of other biochemical and hemogram values (Table 2).

**Table 1.** General characteristics of the study groups.

Baseline characteristics	Diabetic patients with CAE (n=124)	Diabetic patients with NCA (n=124)	P value
Age (years)	61.4 $\pm$ 10.9	59.3 $\pm$ 10.3	0.134
Male/female	70/54	42/82	0.447
LVEF (%)	58.48 $\pm$ 4.34	59.07 $\pm$ 4.61	0.297
Heart rate	73,5 (50-100)	76 (57-107)	0.352
SBP (mmHg)	120 (100-150)	120 (90-158)	0.577
DBP (mmHg)	70 (63-94)	80 (60-100)	0.210
HbA1c (%)	7.3 (4.4-12.0)	6.9 (5.6-12.8)	0.489
BMI	32.0 (18.7-41.1)	31.2 (20.1-41.9)	0.628

CAE: Coronary artery ectasia, NCA: Normal coronary artery, SBP: Systolic blood pressure, DBP: Diastolic blood pressure, HbA1c (%): Hemoglobin A1c, BMI: Body mass index.

**Table 2.** Laboratory data of the study groups.

Parameters	Diabetic patients with CAE (n=124)	Diabetic patients with NCA (n=124)	<i>p</i>
LDL-cholesterol (mg/dL)	112 (49-271)	122 (46-211)	0.336
Triglyceride (mg/dL)	162 (75-631)	166 (70-548)	0.076
Total cholesterol (mg/dL)	189 (126-586)	208 (95-294)	0.637
HDL-cholesterol (mg/dL)	44 (24-75)	46 (29-86)	0.392
Glomerular filtration rate (%)	84 (32-110)	87 (42-125)	0.573
ALT (u/l)	17 (9-50)	20 (9-132)	0.277
AST (u/l)	19 (12-42)	21 (7-48)	0.256
TSH	1.4 (0.3-4.5)	1.5 (0.3-4.2)	0.054
CRP (mg/L)	1 (0.10-18)	1 (0.01-11.2)	0.012
WBC, (u/mm <sup>3</sup> )	8 (4-13)	7 (5-12)	0.023
Hemoglobin (gr/dL)	13 (10-16)	14 (10-20)	0.015
MCV	86 (64-99)	87 (79-97)	0.918
RDW (%)	16 (14-20)	15 (12-19)	0.026
Neutrophil, (u/mm <sup>3</sup> )	4.5 (2.1-11.4)	4.0 (0.2-7.5)	0.003
Lymphocyte, (u/mm <sup>3</sup> )	2.2 (1.2-3.2)	2.5 (0.1-3.6)	0.102
Monocyte, (u/mm <sup>3</sup> )	0.5 (0.2-1)	0,5 (0.2-4.3)	0.589
Basophils, (u/mm <sup>3</sup> )	0.06 (0.001-0.1)	0.07 (0.001-0.2)	0.888
Eosinophil, (u/mm <sup>3</sup> )	0.143 (0.009-0.658)	0.138 (0.033-0.815)	0.846
Platelet counts (Plt) (k/mm <sup>3</sup> )	266 (196-450)	236 (163-362)	0.001
PDW (%)	17.9 (16.2-20.4)	17.7 (15.9-19.7)	0.011
MPV (fL)	8.4 (6.4-11.2)	7.9 (6.6-10.1)	0.015
PCT	0.20 (0.14-0.32)	0.19 (0.13-0.26)	0.001
Neutrophil lymphocyte ratio (NLR)	2.1 (1.0-9.7)	1.6 (0.2-5.7)	0.002
Platelet lymphocyte rate (PLR)	132.5 (71.2-371.9)	97.6 (64.8-4085.7)	0.516

CAE: Coronary artery ectasia, NCA: Normal coronary artery, GFR: glomerular filtration rate, ALT: alanine aminotransferase, AST: aspartate aminotransferase, PDW: Platelet distribution width; RDW: Red cell distribution width; MPV: Mean platelet volume; HDL: high-density lipoprotein; LDL: low-density lipoprotein; WBC: White blood count; PCT: plateletcrit.

## Discussion

This study showed that hemogram parameter levels were different between CAE patients and controls. To the best of our knowledge, this is

the first study to evaluate hemogram parameters in patients with diabetic coronary artery ectasia.

CAE has been related to rising morbidity and

mortality [15]. Angiographies performed to investigate ischemic heart disease indicate an average of 1-5% CAE [4]. CAE is thought an atypical variant of coronary atherosclerosis, that characterized by disruption of the elastic lamina [4, 5]. The key role of inflammation in the initiation and progression of atherosclerosis is well known [22, 23].

Inflammation plays a major role in the development of atherosclerosis and in all stages of CAD [24]. Chronic inflammation is considered to play a role in the etiology of CAE [25, 26]. In our study, as in previous studies [27], the level of inflammatory biomarker CRP was found to be high.

Circulating white blood cell count (WBC) and its subtypes and their relationship to cardiovascular outcomes have been evaluated in previous studies [28]. Leukocyte, monocyte, and neutrophil levels have found to be high in patients with isolated CAE [29]. Neutrophil lymphocyte ratio (NLR) is being evaluated as a new marker of inflammation. Recently, it has been suggested that the NLR rate is a new biomarker for cardiovascular events and prognosis [30]. Balta et al [31] investigated the relationship of NLR in isolated CAE patients and found it to be high. In our study, this rate was high in CAE patients.

Increase in Mean platelet volume (MPV) and platelet distribution width (PDW) in diabetic patients is thought to be related to diabetic vascular complications [32]. MPV and PDW levels have been found to be high in diabetic patients with macrovascular complications [33]. MPV and PDW were higher in diabetic patients with thromboembolic complications [34]. This confirms that CAE is an atherosclerotic disease. MPV, an indicator of platelet activation, has an independent effect on the pathophysiology of atherosclerosis. MPV levels were high in patients with acute

myocardial infarction, unstable angina pectoris, and congestive heart failure [35]. As in our study, MPV levels were found to be high in CAE patients in previous studies [36]. In contrast to this study, in our study, all CAE patients were diabetic. In our study, patients with coronary artery ectasia had higher MPV than normal coronary arteries.

Platelets have an important role in the pathogenesis of homeostasis and thrombosis [37]. It has been shown in previous studies that platelet indices are increased in diabetic patients [38]. We also found these indexes high in our study.

### **Conclusions**

Routine hematological analyzes are important, simple, effortless and cost-effective tests. These tests may be predictive of CAE, which requires prospective large-scale randomized control trials.

**Funding:** *There is no financial support and sponsorship*

**Conflict of Interest:** *The authors declare that they have no conflict of interest.*

**Ethical statement:** *The study was conducted in accordance with the ethical approval of the University Ethics Committee. (Date: 24/10/2019; Decision number: 2019/217).*

### **ORCID iD of the author(s)**

*Mehmet Inanir / 0000-0003-1784-3584*

### **References**

- [1] Endoh S, Andoh H, Sonoyama K, et al. Clinical Features of Coronary Artery Ectasia. J Cardiol. 2004; 43(2):45–52.
- [2] Befeler B, Aranda MJ, Embi A, et al. Coronary artery aneurysms: study of the etiology, clinical course and effect on left

- ventricular function and prognosis. *Am J Med.* 1977; 62(4):597–607.
- [3] Zeina AR, Sharif D, Blinder J, et al. Noninvasive assessment of coronary artery ectasia using multidetector computed tomography. *Coron Artery Dis.* 2007; 18(3):175-80.
- [4] Sultana R, Sultana N, Ishaq M, et al. The prevalence and clinical profile of angiographic coronary ectasia. *J Pak Med Assoc.* 2011; 61(4):372-75.
- [5] Demopoulos VP, Olympios CD, Fakiolas CN, et al. The natural history of aneurysmal coronary artery disease. *Heart.* 1997; 78(2):136-41.
- [6] Díaz-Zamudio M, Bacilio-Pérez U, Herrera-Zarza MC, et al. Coronary artery aneurysms and ectasia: role of coronary CT angiography. *Radiographics.* 2009; 29(7):1939-54.
- [7] al-Harathi SS, Nouh MS, Arafa M, et al. Aneurysmal dilatation of the coronary arteries: diagnostic patterns and clinical significance. *Int J Cardiol.* 1991; 30(2):191-94.
- [8] Manginas A, Cokkinos DV. Coronary artery ectasias: imaging, functional assessment and clinical implications. *Eur Heart J.* 2006; 27(9):1026-31.
- [9] Sarli B, Baktir AO, Saglam H, et al. Neutrophil-to-lymphocyte ratio is associated with severity of coronary artery ectasia. *Angiology.* 2014; 65(2):147-51.
- [10] Antoniadis AP, Chatzizisis YS, Giannoglou GD. Pathogenetic mechanisms of coronary ectasia. *Int J Cardiol.* 2008; 130(3):335–43.
- [11] Mavrogeni S. Coronary artery ectasia: from diagnosis to treatment. *Hellenic J Cardiol.* 2010; 51(2):158–63.
- [12] Neunteufl T, Katzenschlager R, Hassan A, et al. Systemic endothelial dysfunction is related to the extent and severity of coronary artery disease. *Atherosclerosis.* 1997; 129(1):111–18.
- [13] Li JJ, He JG, Nan JL, et al. Is systemic inflammation responsible for coronary artery ectasia? *Int J Cardiol.* 2008; 130(2):e69–e70.
- [14] Aydin M, Tekin IO, Dogan SM, et al. The levels of tumor necrosis factor-alpha and interleukin-6 in patients with isolated coronary artery ectasia. *Mediators Inflamm.* 2009; 2009:106145.
- [15] Zografos TA, Korovesis S, Giazitzoglou E, et al. Clinical and angiographic characteristics of patients with coronary artery ectasia. *Int J Cardiol.* 2013; 167(4):1536–41.
- [16] Papadakis MC, Manginas A, Cotileas P, et al. Documentation of slow coronary flow by the TIMI frame count in patients with coronary ectasia. *Am J Cardiol.* 2001; 88(9):1030–32.
- [17] Krüger D, Stierle U, Herrmann G, et al. Exercise-induced myocardial ischemia in isolated coronary artery ectasias and aneurysms ("dilated coronopathy"). *J Am Coll Cardiol.* 1999; 34(5):1461–70.
- [18] American Diabetes Association. 2. Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes-2019. *Diabetes Care.* 2019; 42(Suppl 1):S13–S28.
- [19] Oliveira JEP, Vencio S. Diretrizes da Sociedade Brasileira de Diabetes: 2013-2014/Sociedade Brasileira de Diabetes. São Paulo: AC Farmacêutica. 2014:1-360.
- [20] Endler G, Klimesch A, Sunder-Plassmann H, et al. Mean platelet volume is an

- independent risk factor for myocardial infarction but not for coronary artery disease. *Br J Haematol.* 2002; 117(2):399–404.
- [21] Farias MG, Dal Bó S. The clinical and laboratory importance of mean platelet volume. *J. Bras. Patol. Med. Lab.* 2010; 46(4):275-82.
- [22] Ozde C, Korkmaz A, Kundi H, et al. Relationship between plasma levels of soluble CD40 ligand and the presence and severity of isolated coronary artery ectasia. *Clin Appl Thromb Hemost.* 2018; 24(2):379–86.
- [23] Zhao ZW, Ren YG, Liu J. Low Serum Adropin Levels are Associated with Coronary Slow Flow Phenomenon. *Acta Cardiol Sin.* 2018; 34(4):307–12.
- [24] Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. *N Engl J Med.* 2005; 352(16):1685–95.
- [25] Furtado M, Katzman MA. Examining the role of neuroinflammation in major depression. *Psychiatry Res.* 2015; 229(1-2):27–36.
- [26] Corrado E, Rizzo M, Coppola G, et al. An update on the role of markers of inflammation in atherosclerosis. *J Atheroscler Thromb.* 2010; 17(1):1–11.
- [27] Ozbay Y, Akbulut M, Balin M, et al. The level of hs-CRP in coronary artery ectasia and its response to statin and angiotensin-converting enzyme inhibitor treatment. *Mediators Inflamm.* 2007; 2007:89649.
- [28] Horne BD, Anderson JL, John JM, et al. Which white blood cell subtypes predict increased cardiovascular risk? *J Am Coll Cardiol.* 2005; 45(10):1638–43.
- [29] Kocaman SA, Taçoy G, Sahinarslan A, et al. Relationship between total and differential leukocyte counts and isolated coronary artery ectasia. *Coron Artery Dis.* 2008; 19(5):307–10.
- [30] Guasti L, Dentali F, Castiglioni L, et al. Neutrophils and clinical outcomes in patients with acute coronary syndromes and/or cardiac revascularisation. A systematic review on more than 34,000 subjects. *Thromb Haemost.* 2011; 106(4):591–99.
- [31] Balta S, Demirkol S, Celik T, et al. Association between coronary artery ectasia and neutrophil-lymphocyte ratio. *Angiology.* 2013; 64(8):627–32.
- [32] Jindal S, Gupta S, Gupta R, et al. Platelet indices in diabetes mellitus: indicators of diabetic microvascular complications. *Hematology.* 2011; 16(2):86–89.
- [33] Alhadas KR, Santos SN, Freitas MMS, et al. Are platelet indices useful in the evaluation of type 2 diabetic patients? *J. Bras. Patol. Med. Lab.* 2016; 52(2):96-102.
- [34] Jabeen F, Fawwad A, Rizvi HA, et al. Role of platelet indices, glycemic control and hs-CRP in pathogenesis of vascular complications in type-2 diabetic patients. *Pak J Med Sci.* 2013; 29(1):152-56.
- [35] Huczek Z, Kochman J, Filipiak KJ, et al. Mean platelet volume on admission predicts impaired reperfusion and long-term mortality in acute myocardial infarction treated with primary percutaneous coronary intervention. *J Am Coll Cardiol.* 2005; 19;46(2):284-90.
- [36] Demir S, Avsar MK, Karakaya Z, et al. Increased mean platelet volume is associated

with coronary artery ectasia. *Postepy Kardiol Interwencyjnej*. 2013; 9(3):241–45.

[37] Li S, Zhu CG, Guo YL, et al. The relationship between the plasma PCSK9 levels and platelet indices in patients with stable coronary artery disease. *J Atheroscler Thromb*. 2015; 22(1):76–84.

[38] Inanir M, Gunes Y, Sincer I. Platelet indices in type 1 diabetes mellitus. *Exp Biomed Res*. 2019; 2(2):69-75.